



Solving the two-dimensional CIS problem by a rational algorithm[☆]

Yurii A. Al'pin^a, Alan George^b, Khakim D. Ikramov^{c,*}

^a*Faculty of Mathematics and Mechanics, Kazan State University, 420008 Kazan, Russian Federation*

^b*Department of Computer Science, University of Waterloo, Waterloo, Ont., Canada*

^c*Faculty of Computational Mathematics and Cybernetics, Moscow State University, 119899 Moscow, Russian Federation*

Received 3 November 1999; accepted 29 February 2000

Submitted by G.P. Barker

Abstract

The CIS problem is formulated as follows. Let p be a fixed integer, $1 \leq p < n$. For given $n \times n$ complex matrices A and B , can one verify whether A and B have a common invariant subspace of dimension p by a procedure employing a finite number of arithmetical operations? We describe an algorithm solving the CIS problem for $p = 2$. Unlike the algorithm proposed earlier by the second and third authors, the new algorithm does not impose any restrictions on A and B . Moreover, when A and B generate a semisimple algebra, the algorithm is able to solve the CIS problem for any p , $1 < p < n$. © 2000 Published by Elsevier Science Inc. All rights reserved.

Keywords: Common invariant subspace; Rational algorithm; 2-generated matrix algebra; Radical; Socle; Shemesh's theorem

1. Introduction

Let A and B be $n \times n$ complex matrices. The CIS problem (where CIS is an abbreviation for “Common Invariant Subspace”) is formulated as follows.

[☆] This work was supported by Natural Sciences and Engineering Research Council of Canada grant OGP0008111.

* Corresponding author.

E-mail address: ikramov@cmc.msk.su (K.D. Ikramov).